

## Psyllium hydrophilic mucilloid oral powder for constipation

SESLHDPR/447

### POLICY STATEMENT

The Registered Nurse (RN) / Registered Midwife (RM) is authorised to instigate nurse/midwife initiated medication without an authorised prescriber's order under the specific circumstances set out in the **INDICATIONS** section and provided there are no contraindications present.

It is important for nursing and midwifery staff to remain aware that:

- Minor ailments may be symptoms of other more serious diseases or may be adverse reactions to medication already prescribed
- Nurse-initiated medication may interact with the patient's prescribed medication
- The maximum daily recommended dose of the medication must not be exceeded<sup>1</sup>

The administering nurse/midwife must record the administration on an approved paper or electronic medication chart, clearly indicating that the medicine was nurse initiated.

If the patient continues to require the medication (i.e. more than two doses in 24 hours) then a medical officer (MO) must be consulted and a regular or PRN order obtained.

A change in the patient's condition such as newly occurring or increasing severity of symptoms must be reported to the MO and investigated.

### INDICATIONS

Constipation in patients aged 6 years and over

### CONTRAINDICATIONS

Intestinal obstruction, partial or complete

Hypersensitivity to the active substance or to any of the excipients.

### PRECAUTIONS

Dysphagia—avoid use; oesophageal obstruction may occur.

Phenylketonuria—avoid Metamucil Smooth Texture<sup>®</sup> -it contains aspartame.

Heart failure or renal impairment – risk of fluid overload

Safe to use in pregnancy and breastfeeding<sup>4</sup>

### HISTORY/ASSESSMENT

- Assess patient's usual bowel habits (frequency of stools, volume, colour, consistency)
- Patient's current bowel status (last time bowel opened)
- Assess for alterations in bowel patterns
- Refer to medical officer if patient has the following symptoms: blood in stools, weight loss, abdominal pain
- Assess patient for faecal impaction.
- Review patient's current medication for medicines which may cause constipation
- Consider risk of fluid overload

**Psyllium hydrophilic mucilloid oral powder  
for constipation**

**SESLHDPR/447**

**PROTOCOL/ADMINISTRATION GUIDELINES**

Caution: CHECK for allergies and/or contraindications			
Drug	Dose	Route	Frequency
Psyllium hydrophilic mucilloid oral powder	<b>6 to 11 years:</b> 5.5 g ( = 1.5 teaspoonsful) in 125 mL of water	Oral	Once
	<b>12 years and over:</b> 11 g ( = 3 teaspoonsful) in 250 mL of water		
Mix in water, stir briskly, and administer immediately. If mixture thickens, add more water and stir.			

**MONITORING - POTENTIAL ADVERSE EFFECTS/INTERACTIONS**

Monitor bowel function and complete stool chart

Common Adverse effects (>1%) : flatulence, bloating, abdominal discomfort.

Rare Adverse effects (<0.1%): hypersensitivity reactions including rhinitis, urticaria, bronchospasm and anaphylactic shock, intestinal obstruction, oesophageal obstruction.

Other medications must be given two hours apart from psyllium hydrophilic mucilloid (Metamucil™) to avoid interactions.

**DOCUMENTATION**

A record of the administration must be made on the approved paper or electronic medication chart noting that the medication was nurse initiated.

A further record of the medication administered including indication, dose and effect must be included in the patient’s health care record.

**PRACTICE POINTS**

- Onset of action is 3 days.
- Should not be given immediately before going to bed.
- Ensure adequate fluid intake
- Ensure adequate dietary fibre intake.
- Encourage mobility, where possible.
- Consider review by dietician, if appropriate.

**REFERENCES/FURTHER READING**

1. [PD2013\\_043 Medication Handling in NSW Public Health Facilities](#)
2. [Product Information Metamucil®](#). MIMS online. Accessed 20/02/2018.
3. [Product Information Fibre Health®](#) MIMS Online. Accessed 17/05/18
4. [eTG complete](#). Melbourne: Therapeutic Guidelines Ltd. November 2017. Accessed 20/02/2018.
5. Rossi S. [Australian Medicines Handbook](#). South Australia: Australian Medicines Handbook Pty Ltd, January 2018.

**Psyllium hydrophilic mucilloid oral powder  
for constipation**

**SESLHDPR/447**

**REVISION and APPROVAL HISTORY**

<b>Date</b>	<b>Revision Number</b>	<b>Author and Approval</b>
July 2015	DRAFT	Pharmacy Department, Prince of Wales Hospital
September 2015	1	Approved by SESLHD Drug & QUM Committee
May 2018	Draft 2	Reviewed by nursing and pharmacy staff. References updated, brand names removed
July 2018	2	Approved by SESLHD Quality Use of Medicines Committee.
September 2021	DRAFT 3	Reviewed by nursing and pharmacy staff. Minor wording updates
October 2021	3	Approved by SESLHD Quality Use of Medicines Committee