

Glycerol (glycerin) suppository for constipation

SESLHDPR/454

POLICY STATEMENT

The Registered Nurse (RN) / Registered Midwife (RM) is authorised to instigate nurse/midwife initiated medication without an authorised prescriber's order under the specific circumstances set out in the **INDICATIONS** section and provided there are no contraindications present.

It is important for nursing and midwifery staff to remain aware that:

- Minor ailments may be symptoms of other more serious diseases or may be adverse reactions to medication already prescribed
- Nurse-initiated medication may interact with the patient's prescribed medication
- The maximum daily recommended dose of the medication must not be exceeded¹

The administering nurse/midwife must record the administration on an approved paper or electronic medication chart, clearly indicating that the medicine was nurse initiated.

If the patient continues to require the medication (i.e. more than two doses in 24 hours) then a medical officer (MO) must be consulted and a regular or PRN order obtained.

A change in the patient's condition such as newly occurring or increasing severity of symptoms must be reported to the MO and investigated.

INDICATIONS

Rapid relief of constipation or when oral laxatives are inappropriate.

CONTRAINDICATIONS

Hypersensitivity to the active substance or to any of the excipients.

PRECAUTIONS

Safe in pregnancy and breastfeeding⁴

HISTORY/ASSESSMENT

- Assess patient's usual bowel habits (frequency of stools, volume, colour, consistency)
- Patient's current bowel status (last time bowel opened)
- Assess for alterations in bowel patterns
- Refer to medical officer if patient has the following symptoms: blood in stools, weight loss, abdominal pain
- Assess patient for faecal impaction.
- Review patient's current medication for medicines which may cause constipation

**Glycerol (glycerin) suppository
for constipation**

SESLHDPR/454

PROTOCOL/ADMINISTRATION GUIDELINES

Caution: CHECK for allergies and/or contraindications			
Drug	Dose	Route	Frequency
Glycerol 700 mg INFANT suppository	<i>Infants 1 month – 1 year:</i> ONE 700 mg suppository	Rectal	Once
Glycerol 1.4 g CHILD suppository	<i>Children 1- 12 years:</i> ONE 1.4 g suppository		
Glycerol 2.8 g ADULT suppository	<i>Over 12 years of age:</i> ONE 2.8 g suppository		
Moisten the tip of the suppository with sterile water and insert well up in the rectum with patient lying on their left side. Allow to remain in the rectum for 15 to 30 minutes.			

MONITORING - POTENTIAL ADVERSE EFFECTS/INTERACTIONS

Monitor bowel function and complete stool chart
Adverse effects: rectal discomfort, mucosal erosion

DOCUMENTATION

A record of the administration must be made on the approved paper or electronic medication chart noting that the medication was nurse initiated.

A further record of the medication administered including indication, dose and effect must be included in the patient’s health care record.

PRACTICE POINTS

- Onset of action is five to 30 minutes.
- The suppository does not need to dissolve completely to be effective
- Ensure adequate fluid and dietary fibre intake
- Encourage mobility, where possible
- Consider review by dietician, if appropriate

REFERENCES/FURTHER READING

1. [PD2013_043 Medication Handling in NSW Public Health Facilities](#)
2. [Product Information Glycerol®](#). MIMS online. Accessed 17/05/2018.
3. [eTG complete](#). Melbourne: Therapeutic Guidelines Ltd. March 2016. Accessed 17/05/2018.
4. [Australian Medicines Handbook](#). South Australia: Australian Medicines Handbook Pty Ltd, 2018.
5. [BNFC 2015](#). Accessed via CIAP 17/05/2018.

**Glycerol (glycerin) suppository
for constipation**

SESLHDPR/454

REVISION and APPROVAL HISTORY

Date	Revision Number	Author and Approval
July 2015	DRAFT	Pharmacy Department, Prince of Wales Hospital
September 2015	1	Approved by SESLHD Drug & QUM Committee
May 2018	DRAFT 2	Reviewed by nursing and pharmacy staff. Minor wording updates made. References updated.
July 2018	2	Approved by SESLHD Quality Use of Medicines Committee
September 2021	DRAFT 3	Reviewed by nursing and pharmacy staff.
October 2021	3	Approved by SESLHD Quality Use of Medicines Committee