

# SESLHD PROCEDURE COVER SHEET



**Health**  
South Eastern Sydney  
Local Health District

<b>NAME OF DOCUMENT</b>	Pain Management - Epidural Analgesia (Adult) non-obstetric
<b>TYPE OF DOCUMENT</b>	Procedure
<b>DOCUMENT NUMBER</b>	SESLHDPR/324
<b>DATE OF PUBLICATION</b>	October 2020
<b>RISK RATING</b>	Medium
<b>LEVEL OF EVIDENCE</b>	National Safety and Quality Standard: 4.4 Medication Safety
<b>REVIEW DATE</b>	October 2023
<b>FORMER REFERENCE(S)</b>	SESLHD PD 302 - Pain Management - Epidural Analgesia - Continuous Infusion (Adult)
<b>EXECUTIVE SPONSOR or EXECUTIVE CLINICAL SPONSOR</b>	Dr Greg Keogh SESLHD Clinical Stream Director, Surgery, Perioperative and Anaesthetics
<b>AUTHOR</b>	Matthew Fay, Clinical Nurse Consultant, Pain Management, St George Hospital <a href="mailto:Matthew.Fay@health.nsw.gov.au">Matthew.Fay@health.nsw.gov.au</a> and Bernadette Bugeja, Clinical Nurse Consultant, Pain Management, Prince of Wales Hospital <a href="mailto:Bernadette.Bugeja@health.nsw.gov.au">Bernadette.Bugeja@health.nsw.gov.au</a> in consultation with SESLHD Pain Management and Anaesthetic departments
<b>POSITION RESPONSIBLE FOR THE DOCUMENT</b>	Andrewina Piazza-Davies SESLHD Clinical Stream Nurse Manager, Surgery, Perioperative, Anaesthetics, ED and Trauma Services <a href="mailto:Andrewina.Piazza-Davies@health.nsw.gov.au">Andrewina.Piazza-Davies@health.nsw.gov.au</a>
<b>KEY TERMS</b>	Epidural analgesia, continuous infusion, patient controlled epidural analgesia, programmed intermittent epidural bolus, epidural catheter
<b>SUMMARY</b>	This document outlines the requirements for the safe management of adult (non-obstetric) patients receiving epidural analgesia via a continuous infusion / patient-controlled / programmed intermittent bolus for pain management.

**COMPLIANCE WITH THIS DOCUMENT IS MANDATORY**

**This Procedure is intellectual property of South Eastern Sydney Local Health District.**

**Procedure content cannot be duplicated.**

Feedback about this document can be sent to [SESLHD-ExecutiveServices@health.nsw.gov.au](mailto:SESLHD-ExecutiveServices@health.nsw.gov.au)

## Pain Management – Epidural Analgesia (Adult) non-obstetric

SESLHDPR/324

### 1. POLICY STATEMENT

This procedure relates to the management of adult patients (non-obstetric) receiving epidural analgesia via a continuous infusion, patient controlled or programmed intermittent bolus for pain management. Epidural analgesia is used selectively for certain post-surgical, medical and trauma patients to provide optimum pain relief safely and effectively via the epidural route.

### 2. BACKGROUND

Epidural analgesia is an effective modality of pain management that provides pain relief by continuous, patient controlled or intermittent administration of pharmacological agents, usually local anaesthetic with or without an opioid, into the epidural space via an indwelling catheter.

#### 2.1 Definitions:

- Authorised prescriber: Anaesthetists / Anaesthetic (Consultants, Fellows and Registrars)
- Continuous Infusion: program delivering a continuous rate as a slow infusion across the entire hour
- Programmed Intermittent Epidural Bolus (**PIEB**) 'Intermittent infusion': delivers background hourly dose as one or more mandatory bolus doses, rather than a slow continuous infusion across the entire hour
- Patient Controlled Epidural Analgesia (**PCEA**): a mode whereby patients may manage their pain with a self-administered bolus of additional epidural solution on demand via the PCEA handset.

### 3. RESPONSIBILITIES<sup>1 2</sup>

- Registered Nurses (RNs)
- Medical Staff – Anaesthetists / Anaesthetic (Consultants, Fellows, Registrars).  
ALL decisions relating to the management of Epidural Analgesia (Adult) non-obstetric patients are only to be made by an Anaesthetist / Anaesthetic (Consultants, Fellows and Registrars) or APS.  
All other medical staff must request Anaesthetic or APS review of patient.

### 4. PROCEDURE

#### 4.1 Education: <sup>1 4</sup>

- All staff involved with the administration of epidural analgesia must have received education / training to manage patients safely and effectively.
- Patients receiving epidural analgesia should only be managed in wards and units where the nursing staff have received education / training in epidural analgesia management.
- Epidural rescue bolus doses may be administered only by RNs who have received education / training in their individual hospital.

#### 4.2 Prescription<sup>2 3 5 6 7 8 9 10 22</sup>

- Epidural analgesia must be prescribed on the approved [NSW Health Epidural Analgesia Adult form SMR130.022](#) in accordance with [NSW Ministry of Health Policy Directive PD2013\\_043 - Medication Handling in NSW Public Health Facilities](#). The infusion must be clearly labelled as '**Epidural**' infusion<sup>3</sup>

## Pain Management – Epidural Analgesia (Adult) non-obstetric

SESLHDPR/324

- Epidural analgesia may only be prescribed by an ‘Authorised prescriber’
- Commonly prescribed (preloaded) local anaesthetic agents:
  - 0.2% Ropivacaine (2mg/mL)
  - 0.2% Ropivacaine (2mg/mL) with Fentanyl 2 microg/mL
  - Fentanyl 2microg/mL with Adrenaline (Epinephrine) 2 microg/mL and 0.1% Bupivacaine in 500mL (POWH only: refer to [Prescribing Protocol SESLHDPR/637](#)).

### Epidural analgesia may be delivered by:

(As per local hospital approved practices)

- **Continuous** infusion +/- Rescue bolus dose
- Patient Controlled Epidural Analgesia (**PCEA**)
- Programmed Intermittent Epidural Bolus (**PIEB**)
- Programmed Intermittent Epidural Bolus (**PIEB**) + Patient Controlled Epidural Analgesia (**PCEA**)

### Note:

- Each individual patient’s risk / benefit assessment for epidural analgesia will have been considered by the individual anaesthetist during time of review
- All patients with epidural analgesia must have intravenous access at all times and continued for a minimum of six hours post epidural catheter removal
- Naloxone, Metaraminol and Atropine should be provided and used by the Rapid Response or Code Blue team for the management of severe adverse effects  
Therapeutic anticoagulants **MUST NOT** be given without consent from APS / Anaesthetist.

### 4.3 Preparation of Epidural Infusion<sup>3</sup>

- Use only pre-loaded epidural infusion flasks to help reduce medication error and assist infection control
- Use only dedicated epidural administration sets that are yellow in colour and portless and must be clearly labelled as ‘Epidural infusion’<sup>2</sup>
- Do NOT inject any other medications into the epidural line.

### 4.4 Programming of Epidural Pump<sup>1</sup>

- Only use dedicated infusion pumps with programmed safety limits for epidural analgesia that are easily distinguishable from those used for intravenous and other types of infusions within individual hospitals<sup>2,11</sup>
- A pump used for epidural analgesia must be programmed by two Registered Nurses
- A pump used for epidural analgesia must be programmed according to the parameters set by the prescribing Anaesthetist / Anaesthetic (Consultants, Fellows, Registrars) / APS on the approved NSW Health Epidural Analgesia Adult form SMR130.022.
- The epidural pump settings should be checked at commencement of each shift, patient transfer, prior to administration of a rescue bolus dose and when the flask is changed.

## Pain Management – Epidural Analgesia (Adult) non-obstetric

SESLHDPR/324

### 4.5 Observations<sup>1</sup>

The RN assigned to the patient receiving epidural analgesia is responsible for ensuring the following observations are carried out and documented on the inside of the NSW Health Epidural Analgesia Adult form SMR130.022. The RN must check the patient's observations are between the flags and refer to the management guidelines on the front of the NSW Health Epidural Analgesia Adult form SMR130.022, if observations fall outside these parameters. If in doubt Anaesthetist / Anaesthetics (Consultants, Fellows, Registrar) / APS should be contacted to review the patient.

OBSERVATIONS	FREQUENCY
<b>Pain Assessment Score at rest “R” and with relevant movement “M”</b> <b>Sedation Score</b> <b>Respiratory Rate</b>  <b>Oxygen therapy</b> <b>SpO<sub>2</sub></b>  <b>Blood Pressure</b> <b>Heart Rate</b>	Hourly for the first six (6) hours post epidural insertion / commencement or change in program.  Then 2 <sup>nd</sup> hourly or more frequently if patients clinical condition warrants  Before any rescue bolus dose, and AFTER A RESCUE BOLUS every ten (10) minutes for 30 minutes and then one (1) hour post rescue bolus dose (or more frequently if directed by an anaesthetist).
<b>Infusion rate (mL/hr)</b> <i>Document cumulative infused total (mL)</i>	Hourly for the first six (6) hours, then 2 <sup>nd</sup> hourly (if no change in rate)
<b>Temperature</b>	Every four (4) hours or more frequently if clinically required
<b>Motor Block Assessment</b> Use Bromage scale <i>(See Appendix B)</i>	Every four (4) hours and prior to mobilisation
<b>Sensory Block Assessment (Dermatome level check)</b> <i>(Refer to 4.5.1 for sensory testing instructions)</i>	Every four (4) hours and prior to / or after administration of a rescue bolus dose or more frequently as specified by Anaesthetist / Anaesthetic Registrar / APS.
<b>Epidural catheter insertion site check</b>	Every eight (8) hours – At shift change, check for: Catheter position, integrity of epidural dressing and signs of leakage. Assess for any tenderness, redness or back pain at the epidural site
<b>Bladder function check</b>	Once per shift
<b>Epidural program check with prescription</b>	At shift handover; when flask is changed; program alterations and on patient transfer between wards / units or procedures

**Pain Management – Epidural Analgesia  
(Adult) non-obstetric**

**SESLHDPR/324**

**4.5.1 Sensory Testing (Dermatome Level Check)<sup>1</sup>** (See Appendix B)

- Place ice on an area well away from the possible dermatome cover (e.g. face or forearm) and ask them to tell you how cold it feels to them.
- Apply ice to an area likely to be blocked on the same side of the body and ask the patient "Does this feel the same cold as your face / arm or different?"
- Apply ice to areas above and below this point until it is clear at which level the top and the bottom of the block is.
- Repeat the procedure on the opposite side of the body (Note: a block may be uneven or unilateral).
- Document the blocked dermatomes on the NSW Health Epidural Analgesia Adult form SMR130.022 in the 'Dermatome Level Check section'.
- Record both the upper and lower limits of the block:  
e.g. T7-L1 L=R or R: T7-L1 L: T10-L2

**4.6 Changing the Infusion Bag<sup>1 2</sup>**

- An infusion bag must be checked by two RNs and changed by an RN who has received epidural analgesia education / training in their individual hospital. It must be recorded on the NSW Health Epidural Analgesia Adult form SMR130.022 in the 'Record of epidural administration' section and signed by both RNs. Both RNs must witness the discarded amount and record in the 'Record of epidural solution discarded' section and sign.
- The dedicated epidural giving set must not be changed without consultation with the Anaesthetist/ Anaesthetic (Consultants/ Fellows/ Registrars) / APS. Routine changing of the giving set is NOT required.
- An aseptic no-touch technique must be used when changing medication bags.
- Epidural bags are to be changed once the infusion has run through, with a maximum hang time of 72 hours, unless specified otherwise in the manufacturer's product information (available on MIMS).

**4.7 Administration of a Rescue Bolus Dose<sup>1 11 12</sup>**

A rescue bolus dose should be administered, as prescribed, when a patient is experiencing inadequate analgesia. Administration of multiple rescue boluses is NOT recommended without review by the Anaesthetist / Anaesthetic (Consultants, Fellows, Registrars) / APS.

- Prior to administration check the epidural infusion delivery device and administration set for faults, kinks or disconnection and perform a full set of observations including motor and sensory level of block assessments
- Check catheter insertion site for catheter position or signs of leakage
- Identify a prescription for RESCUE BOLUS DOSE on the NSW Health Epidural Analgesia Adult form SMR130.022
- Check if a rescue bolus has not been given recently
- RNs who have received epidural analgesia education / training in their individual hospital may administer a rescue bolus dose and increase the rate. The dose must be checked and witnessed by a second RN
- Give prescribed rescue bolus dose and increase rate by 1 to 2 mL/hour
- The RN and witnessing RN must record and initial any rescue bolus dose in the 'Epidural Delivery' section on the NSW Health Epidural Analgesia Adult form

## Pain Management – Epidural Analgesia (Adult) non-obstetric

SESLHDPR/324

SMR130.022

- Perform observations after the rescue bolus dose - as per section 4.5
- If pain persists and observations are stable give another rescue bolus dose if at the minimum interval between rescue doses (hours or minutes) as charted on the approved NSW Health Epidural Analgesia Adult form SMR130.022
- Increase the infusion rate in accordance with the prescription *if the maximum rate has not been reached* and if not contra-indicated by sensory testing and/or motor block
- Continue to monitor as per section 4.5
- If pain continues to persist, contact the APS or if after hours contact the Duty Anaesthetist
- If complications occur see section 4.8.

### 4.8 **Management of Epidural Complications**<sup>1 11 13</sup>

On the NSW Health Epidural Analgesia Adult form SMR130.022 the observations of sedation, respiratory rate, blood pressure, heart rate and motor blockade are colour coded to provide an alert for a Yellow zone (Clinical Review) or Red zone (Rapid Response or Code Blue).

#### 4.8.1 **Respiratory Depression / Over sedation**

Concurrent use of parenteral opioids and sedatives increase the risk of respiratory depression. No other opioids or sedatives should be prescribed unless ordered by Anaesthetist / Anaesthetic (Consultants, Fellows, Registrars) / APS.

#### **If respiratory rate < 10 and / or increasing sedation (Sedation score 2)**

- Ensure oxygen therapy in progress and support airway if necessary
- Encourage patient to breathe deeply
- Activate a Clinical review
- Contact Anaesthetist / Anaesthetic (Consultants, Fellows, Registrars) / APS

#### **If respiratory rate $\leq$ 5 or patient is unrousable (sedation score 3):**

- Stop infusion
- Give oxygen at 15 Litre/minute and support airway if necessary
- Activate a Code Blue if respiratory arrest appears likely
- Administer IV Naloxone if prescribed.

#### 4.8.2 **Hypotension**

Sympathetic blockade may lead to hypotension. However, with low concentrations of anaesthetic drugs used for epidural infusions, hypotension may be the result of hypovolemia rather than the epidural infusion. Other causes of hypotension must always be investigated such as bleeding, sepsis, myocardial insufficiency, pulmonary embolus and dehydration.

## Pain Management – Epidural Analgesia (Adult) non-obstetric

SESLHDPR/324

If systolic is <90mmHg or as adjusted on the Alterations to Calling Criteria on the Electronic Observation Chart (BTF) via eMR.

- Stop the infusion
- Lie patient flat and do not lie head down
- Contact Anaesthetist/ Anaesthetic (Consultants, Fellows, Registrars) / APS
- Activate a Rapid Response review
- Consider IV fluid bolus
- Consider vasopressor

### 4.8.3 Bradycardia

If heart rate <40 or as adjusted on the Alterations to Calling Criteria blood on the Electronic Observation Chart (BTF) via eMR.

- Stop epidural infusion
- Activate Rapid Response
- Atropine must be available in the clinical area
- Contact Anaesthetist/ Anaesthetic (Consultants, Fellows, Registrars) / APS

### 4.8.4 Motor Block Assessment

- If Bromage score 1, 2 or 3 DO NOT ambulate
- Activate a Clinical review
- If patient experiences severe back pain, increasing motor block, bladder / bowel incontinence, numbness / tingling and weakness in lower limbs, contact Anaesthetist / Anaesthetic (Consultants, Fellows, Registrars) / APS **urgently**. The presence of these observations must be reported to a consultant anaesthetist.

### 4.8.5 Inadequate Analgesia<sup>1</sup>

- Give prescribed epidural rescue bolus dose and increase rate within prescription limits – as per section 4.7
- If required, repeat as long as the minimum interval between rescue doses (hours or minutes) as charted on the approved NSW Health Epidural Analgesia Adult form SMR130.022 has elapsed.
- If analgesia is inadequate after second bolus, notify Anaesthetist / Anaesthetic (Consultants, Fellows, Registrars) / APS

### 4.8.6 Nausea or Vomiting

- Administer PRN antiemetics as prescribed on the patient's Medication Administration Record (MAR) via Electronic Medical Record system (eMR)
- If adverse effect continues contact APS.

### 4.8.7 Urinary Retention

- Bladder scan
- Contact patient's primary care team for assessment ± urinary catheterisation

### 4.8.8 Pruritus

- Notify Anaesthetist / Anaesthetic (Consultants, Fellows, Registrars) / APS
- Consider low dose naloxone
- DO NOT give sedating antihistamine

## Pain Management – Epidural Analgesia (Adult) non-obstetric

SESLHDPR/324

### 4.8.9 Inadvertent Epidural Catheter Disconnection

- If catheter is disconnected at the filter, **DO NOT** reconnect
- Stop the infusion
- Cover the catheter end with sterile cap or sterile gauze if catheter disconnects from connector / filter
- Contact Anaesthetist / Anaesthetic (Consultants, Fellows, Registrars) / APS immediately.

### 4.8.10 Epidural Dressing Detaching / Lifting

- Reinforce only if catheter insertion site is not exposed / do not change dressing routinely (use only vapour permeable dressings)
- If insertion site exposed contact Anaesthetist / Anaesthetic (Consultants, Fellows, Registrars) / APS.

## 4.9 Potential Serious Epidural Complications

### 4.9.1 Post Dural Puncture Headache<sup>14</sup>

Signs and symptoms: headache (bifrontal or occipital) are usually postural (exacerbated when patient is in an upright position and improved when lying flat)

- Contact Anaesthetist / Anaesthetic (Consultants, Fellows, Registrars) / APS
- Treatment includes lying flat, bed rest, analgesia (simple or opioid), high fluid intake (unless contraindicated) and caffeine
- Consider epidural blood patch.

### 4.9.2 Epidural Haematoma<sup>15</sup>

The puncture of epidural blood vessels during catheter insertion or removal may result in the formation of an epidural haematoma particularly in the presence of coagulopathy

Signs and symptoms:

- lower limb weakness and / or numbness (increasing motor block)
- back pain
- bowel or bladder dysfunction
- Stop infusion, call APS / Anaesthetist / medical team for **immediate urgent neurological assessment** / may need urgent MRI / urgent surgical decompression if neurological changes develop due to nerve or spinal cord compression.

### 4.9.3 Epidural Space Infection

- May be prevented by using strict aseptic technique during insertion, preparation and administration of solutions. Always connect epidural line to a bacterial filter, ensure all connections are Luer Locked, secured and maintain a clear occlusive dressing over epidural site
- If patient has temperature spikes > 38.5°C, notify APS / Anaesthetics and consider removal of epidural catheter
- If signs of inflammation / infection at insertion site, notify APS, consider removal of epidural catheter
- The presence of severe or increasing back pain, may indicate epidural abscess and should be investigated promptly (even in the absence of fever) see 4.9.2
- As epidural space infection can present up to six weeks post epidural catheter



## Pain Management – Epidural Analgesia (Adult) non-obstetric

SESLHDPR/324

removal, patients should be educated regarding signs and symptoms and written post epidural information should be provided where available

### 4.9.4 Neurological Injury<sup>15</sup>

Direct damage to the spinal cord or peripheral nerves due to the epidural needle or catheter is extremely rare.

- Signs and symptoms - weakness, numbness, tingling sensation in lower limbs, bowel or bladder incontinence
- Stop infusion, call APS / Anaesthetist / medical team for **immediate urgent neurological assessment**.

### 4.9.5 Catheter Migration<sup>16</sup>

Rarely a catheter placed in the epidural space may migrate into the intrathecal space or an epidural blood vessel.

- Signs and symptoms - migration into the intrathecal space will usually result in a rapidly increasing block with a sudden onset of complications. Migration into a blood vessel may result in increasing pain +/- signs of local anaesthetic toxicity (perioral numbness, tinnitus, dizziness, facial twitching, seizures)
- Stop infusion, notify APS / medical team and activate a clinical review or rapid response according to observations.

## 5. CONCURRENT ANTICOAGULANT MEDICATIONS<sup>17 18 19 20 21</sup>

- Each individual patient's risk / benefit assessment for insertion of an epidural needle / catheter is the responsibility of the anaesthetist
- Anticoagulation is the most important risk factor for the development of epidural haematoma following insertion of epidural needle / catheter
- **Therapeutic anticoagulation is contraindicated whilst epidural catheters remain in situ, however most patients do receive anticoagulant medications for thromboprophylaxis**
- It is vital that adequate time delays exist between the administration of anticoagulants and the removal of epidural catheters

### 5.1 Removal of Epidural Catheter<sup>17 18 19 20 21</sup>

An epidural catheter may be ceased and removed only when APS / Anaesthetist has instructed its removal and:

- The patient has **NOT RECEIVED unfractionated subcutaneous Heparin:**
  - **≤5000 IU** dose within the previous **six (6) hours**
  - **>5000 IU** dose within the previous **twelve (12) hours**
- The patient has **NOT RECEIVED a prophylactic dose of low molecular weight heparin (LMWH) e.g. dalteparin or enoxaparin** within the previous **twelve (12) hours**.
- The patient has **NOT RECEIVED** therapeutic dose of:
  - **Rivaroxaban** within the previous **twenty-six (26) hours**
  - **Apixaban** within the previous **thirty (30) hours**
  - **Dabigatran** within the previous **thirty-six (36) hours**
- The patient on warfarin has a documented INR of **<1.5**
- The anaesthetist must leave clear instructions about catheter removal and administration of any anticoagulants that are not discussed in this document
- **No prophylactic doses of anticoagulants are to be administered for at least**

## Pain Management – Epidural Analgesia (Adult) non-obstetric

SESLHDPR/324

**four (4) hours** following the removal of an epidural catheter unless specified by APS / Anaesthetist / Anaesthetic (Consultants, Fellows, Registrar).

### 52 Procedure for Removal of Epidural Catheter<sup>1 17 18 19 20 21</sup>

- Ensure patient's coagulation results are most recent and adhere to 5.1
- Explain the procedure to the patient reassuring them that removal of the catheter is not uncomfortable
- Position patient lying on side or sitting up with spine slightly flexed forward
- Wash hands and organise equipment
- Stop infusion, **do not** wean infusion
- Carefully remove epidural catheter dressing
- Wash hands and put on sterile gloves
- Gently withdraw catheter. Do not forcefully pull out catheter. Contact the Anaesthetist / Anaesthetic (Consultants, Fellow, Registrars) / APS if any resistance is felt when trying to remove catheter
- If signs of infection (purulent drainage, redness or swelling) are present send the epidural catheter tip for culture and notify Anaesthetist / Anaesthetic (Consultants, Fellow, Registrars) / APS
- Cover site with a clear occlusive dressing
- Confirm that epidural catheter tip is intact with second RN and document and sign on the bottom of page 3 of the NSW Health Epidural Analgesia Adult form SMR130.022
- Monitor patient's sensory and motor function every two (2) hours for first six (6) hours, then every four (4) hours for next eighteen (18) hours post epidural catheter removal
- Check the epidural site every eight (8) hours, for twenty-four (24) hours, then at 48 hours for any signs of infections e.g. redness, swelling, discharge, bruising, tenderness or pain<sup>4,15</sup>

### 6. DOCUMENTATION<sup>1</sup>

NSW Health Epidural Analgesia Adult form SMR130.022  
Medical Record  
Approved electronic medication management system

### 7. AUDIT

Patients receiving epidural analgesia will be reviewed daily by Pain Management clinicians and Anaesthetists.

Review of ims+ reports.

### 8. REFERENCES

1. [NSW Health Epidural Analgesia \(Adult\) form SMR130.022](#)
2. [NSW Ministry of Health Policy Directive PD2013\\_043 - Medication Handling in NSW Public Health Facilities](#)
3. [NSW Ministry of Health Policy Directive PD2016\\_058 - User-applied Labelling of Injectable Medicines, Fluids and Lines](#)
4. Macintyre, PE & Schug, SA 2015, *Acute pain management: a practical guide*, 4<sup>th</sup> edn, CRC Press: Boca Raton, Florida.

**Pain Management – Epidural Analgesia  
(Adult) non-obstetric**

SESLHDPR/324

5. Al-Rawi, OY, Pennefather, SH, Page, RD, Dave, I & Russell, G 2008, 'The effect of thoracic epidural bupivacaine and an intravenous adrenaline infusion of gastric tube blood flow during oesophagectomy', *Anesthesia and Analgesia*, vol. 106, no. 3, pp. 884-887, <<https://pubmed.ncbi.nlm.nih.gov/18292435/>>.
6. Fegley, AJ, Lerman, J & Wissler, R 2008, 'Epidural multi-orifice catheters functions as a single-orifice catheters: an in vitro study', *Anesthesia and Analgesia*, vol. 107, no. 3, pp. 1079-1081, <<https://pubmed.ncbi.nlm.nih.gov/18713933/>>.
7. Kuthalia, G & Chaudhary, G 2011, 'Ropivacaine: a review of its pharmacology and clinical use', *Indian Journal of Anaesthesia*, vol. 55, no. 2, pp. 104-110, <[http://www.ijaweb.org/temp/IndianJAnaesth552104-3800163\\_103321.pdf](http://www.ijaweb.org/temp/IndianJAnaesth552104-3800163_103321.pdf)>.
8. Lim, Y, Chakravarty, S, Ocampo, CE & Sia, AT 2010, 'Comparison of automated intermittent low volume bolus with continuous infusion for labour epidural analgesia', *Anaesthesia and Intensive Care*, vol. 38, no. 5, pp. 894-899, <<https://journals.sagepub.com/doi/pdf/10.1177/0310057X1003800514>>.
9. Sng, BL, Kwok, SC & Sia, ATH 2015, 'Modern neuraxial labour analgesia', *Current Opinion in Anaesthesiology*, vol. 28, no. 3, pp. 285-289, <<https://pubmed.ncbi.nlm.nih.gov/25827278/>>.
10. Wong, CA, McCarty, RJ & Hewlett, B 2011, 'The effect of manipulation of the programmed intermittent bolus time interval and injection volume on total drug use for labour epidural analgesia: a randomized controlled trial', *Anesthesia and Analgesia*, vol. 112, no. 4, pp. 904-911, <<https://pubmed.ncbi.nlm.nih.gov/21430035/>>.
11. NHS National patient safety agency 2007, *Patient safety alert: safer practice with epidural injections and infusions*, United Kingdom, <<https://www.sps.nhs.uk/wp-content/uploads/2018/02/2007-NRLS-0396-Epidural-injectns-PSA-2007-03-28-v1.pdf>>.
12. Schug, SA, Palmer, GM, Scott, DA, Halliwell, R & Trinca, J 2015, *Acute Pain Management: Scientific Evidence* 4th edn, Working Group of the Australian and New Zealand College of Anaesthetists and Faculty of Pain Medicine, ANZCA & FPM, Melbourne, <<https://www.anzca.edu.au/getattachment/4c3b03b7-52bf-4c10-9115-83d827c0fc38/Acute-Pain-Management-Scientific-Evidence.aspx>>.
13. Hadzic, A 2019, *Hadzic's Textbook of Regional Anesthesia and Acute Pain Management* 2<sup>nd</sup> edn, McGraw-Hill Education, USA.
14. Obstetric Anaesthetists' Association 2018, 'Treatment of obstetric post-dural puncture headache', Guideline, December 2018, viewed 7 August 2020, <<https://www.oaa-anaes.ac.uk/assets/managed/cms/files/Guidelines/New%20PDPH%20Guidelines.pdf>>.
15. Breivik, H 1998, 'Neurological complications in association with spinal and epidural analgesia', *Acta Anaesthesiologica Scandinavica*, vol. 42, no. 6, pp. 609-613, <<https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1399-6576.1998.tb05290.x>>.
16. [Therapeutic Goods Administration Australian Government 2010, Australian Public Assessment Report for Ropivacaine.](#)
17. Gogarten, W, Vandermeulen, E, van Aken, H, Hozek, S, Llau, JV, & Samama, CM 2010, Royal anaesthesia and antithrombotic agents: recommendations of the European Society of Anaesthesiology, *Eur J Anaesthesio*, vol. 27, pp. 999-1015, <<https://pubmed.ncbi.nlm.nih.gov/20890208/>>.
18. Horlocker, TT, Vandermeulen, E, Kopp, SL, Gogarten, W, Leffert, LR & Benzon, HT 2018, 'Regional anesthesia in the patient receiving antithrombotic or thrombolytic therapy: American society of regional anesthesia and pain medicine evidence-based guidelines' 4th edn, *Regional Anesthesia and Pain Medicine*, vol. 43, no. 3, pp. 263-309, <<https://rapm.bmj.com/content/rapm/43/3/263.full.pdf>>.
19. Horlocker, T 2011, 'Regional anaesthesia in the patient receiving antithrombotic and antiplatelet therapy', *British Journal of Anaesthesia*, vol. 107, Suppl. s1, i96-i106, <<https://pubmed.ncbi.nlm.nih.gov/22156275/>>.
20. The Royal College of Anaesthetists, Royal College of Nursing, The Association of Anaesthetists of Great Britain and Ireland, The British Pain Society and The European

**Pain Management – Epidural Analgesia  
(Adult) non-obstetric**

**SESLHDPR/324**

Society of Regional Anaesthesia and Pain Therapy 2010, *Best practice in the management of continuous epidural analgesia in the hospital in the hospital setting*, The Royal College of Anaesthetists, London, <<https://www.euroespa.com/wp-content/uploads/2014/10/EpiduralAnalgesia2010.pdf>>.

21. Clinical Excellence Commission 2018, 'Guidelines on preoperative management of anticoagulant and antiplatelet agents', Government guideline, December 2018, viewed 7 August 2020, <[http://www.cec.health.nsw.gov.au/\\_data/assets/pdf\\_file/0006/458988/Guidelines-on-perioperative-management-of-anticoagulant-and-antiplatelet-agents.pdf](http://www.cec.health.nsw.gov.au/_data/assets/pdf_file/0006/458988/Guidelines-on-perioperative-management-of-anticoagulant-and-antiplatelet-agents.pdf)>.
22. [Prescribing Protocol SESLHDPR/637 Fentanyl 2microg/mL with Adrenaline \(epinephrine\) 2 microg/mL and Bupivacaine 0.1%.](#)

**9. REVISION AND APPROVAL HISTORY**

Date	Revision No.	Author and Approval
September 2016	1	Reviewed by Drug and Quality Use of Medicine Committee
October 2016	1	Reviewed and approved by Drug and Quality Use of Medicine Committee
August 2020	2	Minor Review. Revised by Matthew Fay, CNC Pain Management, SGH and Bernadette Bugeja, CNC Pain Management, POWH. Document name change; Added section 2.1: definitions and added appendices. Update to sections 4.2, 5.1 and references. Approved by Executive Sponsor.
September 2020	2	Formatted by Executive Services. To be tabled at October Quality Use of Medicines Committee.
October 2020	2	Approved by the Quality Use of Medicines Committee. Published by Executive Services.

**Pain Management – Epidural Analgesia  
(Adult) non-obstetric**

SESLHDPR/324

**Appendix A: Front page – (Management Guidelines & Managing Adverse Effects)**

A copy of the NSW health Epidural chart can be found at: [NSW Health Epidural Analgesia Adult form SMR130.022](https://www.health.nsw.gov.au/epidural/Pages/NSW-Health-Epidural-Analgesia-Adult-form-SMR130.022.aspx)

SMR130022  Holes Punched as per AS2828.1: 2012 BINDING MARGIN - NO WRITING		FAMILY NAME	MRN
	Facility:	GIVEN NAME	<input type="checkbox"/> MALE <input type="checkbox"/> FEMALE
		D.O.B. ____/____/____	M.O.
		ADDRESS	
	<b>EPIDURAL ANALGESIA (ADULT)</b> (Not for use in labour)	LOCATION	
COMPLETE ALL DETAILS OR AFFIX PATIENT LABEL HERE			
<p><b>Epidural Analgesia Management Guidelines</b></p> <p>(For detailed information regarding epidural prescribing and management refer to local hospital policy)</p> <ul style="list-style-type: none"> <li>• <b>Observations</b> on this form to be recorded hourly for 6 hours, then second hourly or more frequently if patient's clinical condition warrants.</li> <li>• <b>Observations AFTER A RESCUE BOLUS</b> (blood pressure and pulse) every 10 minutes for 30 minutes and then 1 hour post bolus (or more frequently if directed by an anaesthetist).</li> <li>• <b>Motor block assessment</b> every four hours and prior to mobilisation.</li> <li>• <b>Dermatome level check</b> refer to local hospital policy.</li> <li>• <b>Catheter site check</b> every 8 hours.</li> <li>• <b>The infusion pump settings</b> to be checked at the commencement of each shift, on patient transfer and when the syringe or bag is changed.</li> <li>• <b>Intravenous access</b> to be maintained for duration of epidural infusion or PCEA.</li> <li>• <b>A dedicated giving set</b> that is yellow in colour and portless must be used.</li> <li>• <b>No other opioids or sedatives</b> to be administered unless ordered by the Acute Pain Service or equivalent medical officer.</li> <li>• <b>Therapeutic anticoagulants MUST NOT</b> be commenced without prior discussion with the Acute Pain Service or equivalent medical officer.</li> <li>• <b>Inadvertent disconnection of epidural catheter from filter: DO NOT</b> re-connect and contact the Acute Pain Service or equivalent medical officer immediately.</li> </ul> <p style="text-align: center;"><b>Managing Adverse Effects</b></p> <ul style="list-style-type: none"> <li>• <b>Motor block or developing leg weakness, severe back pain or tenderness at epidural site could be signs of an epidural haematoma or epidural abscess:</b> Contact the Acute Pain Service or equivalent medical officer immediately. The presence of these observations must also be reported to a consultant anaesthetist.</li> <li>• <b>Hypotension:</b> Refer to instructions below for management guidelines.</li> <li>• <b>Pruritus or persistent nausea or vomiting:</b> Administer PRN medication as prescribed on the patient's National Inpatient Medication Chart. If adverse effect continues contact the Acute Pain Service or equivalent medical officer.</li> <li>• Antihistamines for pruritus are generally ineffective and may contribute to sedation.</li> <li>• <b>Urinary retention:</b> Contact the patient's surgical or medical team.</li> </ul>			
REFER TO YOUR LOCAL CLINICAL EMERGENCY RESPONSE SYSTEM (CERS) PROTOCOL FOR INSTRUCTIONS ON HOW TO MAKE A CALL TO ESCALATE CARE FOR YOUR PATIENT			
<b>APPROPRIATE CLINICAL CARE FOR PATIENTS WITH YELLOW ZONE OR RED ZONE OBSERVATIONS:</b> 1. ENSURE OXYGEN THERAPY IS IN PROGRESS 2. STOP EPIDURAL PUMP FOR ANY RED ZONE OBSERVATIONS 3. ENSURE THAT THE ACUTE PAIN SERVICE OR EQUIVALENT MEDICAL OFFICER IS CONTACTED			
<p><b>YELLOW ZONE RESPONSE</b></p> <p>IF YOUR PATIENT HAS ANY YELLOW ZONE OBSERVATIONS OR additional criteria*                  YOU MUST FOLLOW THE YELLOW ZONE RESPONSE INSTRUCTIONS ON THE NSW STANDARD OBSERVATION CHARTS AND INITIATE APPROPRIATE CLINICAL CARE AS STATED ABOVE</p> <p>*Additional YELLOW ZONE Criteria for Local Anaesthetic Toxicity</p> <ul style="list-style-type: none"> <li>• Numbness and tingling around the mouth and tongue</li> <li>• Metallic taste, tinnitus and dizziness</li> </ul>			
<p><b>RED ZONE RESPONSE</b></p> <p>IF YOUR PATIENT HAS ANY RED ZONE OBSERVATIONS OR additional criteria*                  YOU MUST CALL FOR A RAPID RESPONSE (as per local CERS), FOLLOW THE RED ZONE RESPONSE INSTRUCTIONS ON THE NSW STANDARD OBSERVATION CHARTS AND INITIATE APPROPRIATE CLINICAL CARE AS STATED ABOVE</p> <p>*Additional RED ZONE Criteria for Local Anaesthetic Toxicity</p> <ul style="list-style-type: none"> <li>• Muscular twitching</li> <li>• Convulsion</li> <li>• Cardiovascular collapse</li> </ul>			
<p><b>ACUTE PAIN SERVICE or equivalent medical officer CONTACT:</b></p> <p>BUSINESS HOURS page/phone: _____ OUT OF HOURS page/phone: _____</p>			
NO WRITING <span style="float: right;">Page 1 of 16</span>			

EPIDURAL ANALGESIA (ADULT)  
(Not for use in labour)

SMR130.022

**Pain Management – Epidural Analgesia  
(Adult) non-obstetric**

SESLHDPR/324

**Appendix B: Back page – (Motor and Sensory block assessment)**

A copy of the NSW health Epidural chart can be found at: [NSW Health Epidural Analgesia Adult form SMR130.022](http://www.nsw.gov.au/health/epidural-analgesia-adult-form-smr130.022)

**MOTOR BLOCK ASSESSMENT**

**DO NOT PRINT**

Holes Punched as per AS2928.1:2012  
BINDING MARGIN - NO WRITING

SMR130022