

|                     |  |
|---------------------|--|
| <b>Alert</b>        | <p>Most often given in conjunction with calcium for the prevention and treatment of metabolic bone disease in preterm infants.</p> <p>1 mmol phosphorus/phosphate (P) = 31 mg elemental phosphorus.</p> <p>1 mmol elemental calcium (Ca) = 40 mg elemental calcium.</p> <p>Separate oral doses from calcium supplements by at least 1 hour.</p> <p>When using IV preparation, always check plasma sodium and potassium concentrations to assist in choosing the right phosphate preparation (e.g. sodium or potassium phosphate preparation).</p>  |
| <b>Indication</b>   | <p>Treatment of Metabolic Bone Disease.</p> <p>Treatment of hypophosphataemia.</p> <p>Supplementation to meet the recommended daily intakes.</p>   |
| <b>Action</b>       | Phosphorus is a major intracellular mineral and is important in bone mineralisation and energy production.   |
| <b>Drug Type</b>    | Mineral  |
| <b>Trade Name</b>   | <p><b>IV</b></p> <p>Glycophos® Concentrated injection solution for infusion (Fresenius-Kabi) (recommended organic preparation)</p> <p>Each 1 mL of Glycophos® corresponds to 1 mmol phosphate and 2 mmol sodium.</p> <p>Sodium dihydrogen phosphate Phebra IV (Preferred inorganic preparation)</p> <p>Each 1 mL vial corresponds to 1 mmol phosphate, 1 mmol sodium and 2 mmol hydrogen.</p> <p>Potassium dihydrogen phosphate concentrated injection DBL IV</p> <p>Potassium dihydrogen phosphate concentrated injection Phebra IV</p> <p>Each 1 mL ampoule corresponds to 1 mmol phosphate, 1 mmol potassium and 2 mmol hydrogen.</p> <p><b>ORAL</b></p> <p>Phosphate-Phebra® oral effervescent tablets</p> <p>Each tablet contains: 16.1 mmol phosphate (equivalent to 500 mg elemental phosphorus); 20.4 mmol sodium; 3.1 mmol potassium</p> <p>Sodium dihydrogen phosphate Phebra IV (preferred IV preparation)</p> <p>Each 10 mL vial (sodium dihydrogen phosphate 1.56 g) contains: 10 mmol phosphate; 10 mmol sodium; 20 mmol hydrogen</p> <p>Potassium dihydrogen phosphate concentrated injection DBL IV</p> <p>Potassium dihydrogen phosphate concentrated injection Phebra IV</p> <p>Each 10 mL ampoule (potassium dihydrogen phosphate 1.361 g) contains: 10 mmol phosphate; 10 mmol potassium; 20 mmol hydrogen</p> |
| <b>Presentation</b> | <p><b>IV:</b> Glycophos 20 mL ampoule; Sodium dihydrogen phosphate 10 mL vial; Potassium dihydrogen phosphate concentrated injection 10 mL ampoule.</p> <p><b>Oral:</b> 500 mg effervescent tablets; IV preparation (e.g. sodium or potassium dihydrogen phosphate) can be given orally.</p>   |
| <b>Dose</b>         | <p><b>Treatment of metabolic bone disease (MBD)</b></p> <p>PO: 1 to 3 mmol/kg/day in 2-4 divided doses as an addition to intake from milk and other sources to a maximum intake of 4.5 mmol/kg/day.</p> <p>Use either Sodium dihydrogen phosphate Phebra IV preparation or Phosphate-Sandoz tablets.</p> <p><b>General principles of treatment of MBD:</b></p> <ol style="list-style-type: none"> <li>A. Commence at low dose (e.g. 1 mmol/kg/day) and titrate the dose up as tolerated.</li> <li>B. Given in conjunction with calcium supplementation (but not together - example: <b>Calcium</b> 8 AM, 2 PM, 8 PM and <b>Phosphorus</b> 6 AM, 12 MD, 6 PM )</li> <li>C. Aim to reach the upper end of the recommended intake: Ca 5 mmol/kg/day and P 4.5 mmol/kg/day.<sup>8</sup></li> </ol>   |

|                              |   |
|------------------------------|---|
|                              | <p>D. Dose can be adjusted with a goal of slight excess supply aiming for urinary calcium <math>\geq 1.2</math> mmol/L and phosphate <math>\geq 0.4</math> mmol/L.</p> <p><b>Treatment of acute hypophosphataemia</b><br/>           IV: 0.2 mmol/kg/dose [range 0.15–0.33 mmol/kg/dose] over 6 hours. Repeat as necessary. Aim to maintain normophosphataemia of 1.8–2.6 mmol/L (5.6–8.1 mg/dl).</p> <p><b>Daily enteral Supplementation to meet the recommended daily intakes (RDI)</b><br/>           2–4.5 mmol/kg/day (62–140 mg/kg/day of phosphorus)<sup>7,8</sup></p> <ol style="list-style-type: none"> <li>1. Calculate intake from parenteral and enteral sources</li> <li>2. Supplement the difference via IV or oral route.</li> </ol>   |
| <b>Dose adjustment</b>       |   |
| <b>Maximum dose</b>          |   |
| <b>Total cumulative dose</b> |   |
| <b>Route</b>                 | PO<br>IV  |
| <b>Preparation</b>           | <p><b>IV infusion for treatment of acute hypophosphataemia:</b></p> <p>IV infusion (Glycophos): Draw up 1 mL (1 mmol phosphate) and add 19 mL sodium chloride 0.9% or water for injection to make a final volume of 20 mL with a concentration of 0.05 mmol/mL. Draw up 4 mL/kg (0.2 mmol/kg).</p> <p>IV infusion (sodium dihydrogen phosphate): Draw up 1 mL (1 mmol phosphate) and add 19 mL sodium chloride 0.9% or glucose 5% to make a final volume of 20 mL with a concentration of 0.05 mmol/mL. Draw up 4 mL/kg (0.2 mmol/kg).</p> <p>IV infusion (potassium dihydrogen phosphate): Draw up 1 mL (1 mmol phosphate) and add 24 mL sodium chloride 0.9% or glucose 5% to make a final volume of 25 mL with a concentration of 0.04 mmol/mL. Draw up 5 mL/kg (0.2 mmol/kg).</p> <p><b>Oral</b></p> <p>Option 1 (preferred option for infants going home or when a long storage time is required in the NICU): Disperse 500 mg (16.1 mmol) Phosphate effervescent tablet in 16 mL of water for injection to make a solution with a concentration of 1 mmol/mL.</p> <p>Option 2 (can be used where preparation with low osmolality is preferred e.g. infants with history of feed intolerance): IV sodium dihydrogen phosphate decanted into a bottle and given orally undiluted (expiry time: 7 days).</p> |
| <b>Administration</b>        | <p><b>Oral</b><br/>           Can be administered with feeds (refer to evidence summary section).<br/>           Separate calcium supplements by at least 2 hours.</p> <p><b>IV</b><br/> <b>As part of parenteral nutrition fluid</b> – refer to individual parenteral nutrition formulations.</p> <p><b>IV infusion for treatment of acute hypophosphataemia:</b><br/>           IV glycophos: Infuse over at least 8 hours.<br/>           IV sodium dihydrogen phosphate or IV potassium dihydrogen phosphate: Infuse over at least 6 hours.</p> <p>For severe hypophosphataemia infuse over 8–12 hours. Maximum infusion rate of 0.2 mmol/kg/h.</p>   |
| <b>Monitoring</b>            | <p>Phosphate, calcium, magnesium, alkaline phosphatase concentrations are required at least fortnightly or more often if required. Once these concentrations normalise, serum analysis may be performed once monthly for 6 months or at the discretion of the clinician.<sup>10</sup></p> <p>Urinary calcium and phosphate and Tubular Reabsorption Phosphate (TRP)%, parathormone, and vitamin D concentrations may be useful under certain circumstances .</p>  |

| <b>Contraindications</b>              | Hyperphosphataemia, dehydration, severe renal insufficiency, shock.  |                                   |                                   |                                   |                                       |                 |                 |                 |                   |                  |              |                 |                 |
|---------------------------------------|--|-----------------------------------|-----------------------------------|-----------------------------------|---------------------------------------|-----------------|-----------------|-----------------|-------------------|------------------|--------------|-----------------|-----------------|
| <b>Precautions</b>                    | Hypernatraemia (avoid sodium dihydrogen phosphate).<br>Hyperkalaemia (avoid potassium dihydrogen phosphate)  |                                   |                                   |                                   |                                       |                 |                 |                 |                   |                  |              |                 |                 |
| <b>Drug Interactions</b>              | Calcium and magnesium antacids (e.g. acetate, carbonate, citrate, hydroxide etc.) reduce phosphate absorption — separate doses by at least 2 hours.<br>Additive effects with other drugs that may prolong QT interval.<br>Potassium dihydrogen phosphate preparation may increase the risk of hyperkalaemia when used in conjunction with potassium sparing diuretics (e.g. spironolactone).   |                                   |                                   |                                   |                                       |                 |                 |                 |                   |                  |              |                 |                 |
| <b>Adverse Reactions</b>              | Diarrhoea (oral use only), hypocalcaemia, nephrotoxicity, prolonged QT interval, hypotension, hypomagnesaemia.<br>Hyperphosphataemia – carpedal spasm, seizures. <sup>2</sup>  |                                   |                                   |                                   |                                       |                 |                 |                 |                   |                  |              |                 |                 |
| <b>Compatibility</b>                  | <u>Glycophos</u><br>Fluids: Sodium chloride 0.9%, water for injection, glucose 5%.<br>Y-site: No information.<br><u>Potassium dihydrogen phosphate</u><br>Fluids: Glucose 5%, glucose 10%, glucose in sodium chloride solutions, sodium chloride 0.45%, sodium chloride 0.9%, sodium chloride 3%.<br>Y-site: No information.<br><br><u>Sodium dihydrogen phosphate</u><br>Fluids: Glucose 5%, sodium chloride 0.9%.<br>Y-site: No information  |                                   |                                   |                                   |                                       |                 |                 |                 |                   |                  |              |                 |                 |
| <b>Incompatibility</b>                | <u>Potassium dihydrogen phosphate</u><br>Fluids: No information<br>Drugs: Aciclovir, amiodarone, calcium salts, ketamine, lorazepam, magnesium salts, rocuronium.<br>Solutions that contain other cations such as calcium, magnesium, iron and aluminium may also precipitate.<br><br><u>Sodium dihydrogen phosphate</u><br>Fluids : No information<br>Drugs: Aciclovir, amiodarone, calcium salts, calcium, aluminium or magnesium, iron and magnesium containing solutions.  |                                   |                                   |                                   |                                       |                 |                 |                 |                   |                  |              |                 |                 |
| <b>Stability</b>                      | Preparation from oral effervescent tablets: It is to be used immediately after preparation and discard unused portion.<br>Oral preparation from IV sodium dihydrogen phosphate: 7 days<br>Glycophos: To be used within 24 hours after reconstitution.  |                                   |                                   |                                   |                                       |                 |                 |                 |                   |                  |              |                 |                 |
| <b>Storage</b>                        | Store below 25°C.  |                                   |                                   |                                   |                                       |                 |                 |                 |                   |                  |              |                 |                 |
| <b>Excipients</b>                     | Phosphate-Phebra® oral effervescent tablets: Sodium bicarbonate, potassium bicarbonate, macrogol 4000, citric acid, sucrose, orange 52570 TP0551 and saccharin sodium.<br>Glycophos: Hydrochloric acid and water for injections.   |                                   |                                   |                                   |                                       |                 |                 |                 |                   |                  |              |                 |                 |
| <b>Special Comments</b>               |  |                                   |                                   |                                   |                                       |                 |                 |                 |                   |                  |              |                 |                 |
| <b>Evidence</b>                       | <p><b>Recommended daily intakes (RDI)</b><br/>Phosphorus absorption is typically 80% to 90% of dietary intake.<sup>3</sup></p> <p><b>Parenteral intake:</b> Previously, the recommended doses of <b>parenteral</b> Ca and P in preterm infants varied from 1.3–3 mmol Ca/kg/day and 1.0–2.3 mmol P/kg/day, with a Ca:P ratio in the range of 1.3–1.7.<sup>1,4-6</sup> ESPGHAN 2018 updated guidelines on parenteral nutrition recommends the following Ca and Phosphate:<sup>12</sup></p> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th></th> <th>Parenteral Ca<br/>mmol (mg)/kg/day</th> <th>Parenteral Ph<br/>mmol (mg)/kg/day</th> </tr> </thead> <tbody> <tr> <td>Preterm during the first days of life</td> <td>0.8-2.0 (32-80)</td> <td>1.0-2.0 (31-62)</td> </tr> <tr> <td>Growing preterm</td> <td>1.6-3.5 (100-140)</td> <td>1.6-3.5 (77-108)</td> </tr> <tr> <td>Term neonate</td> <td>0.8-1.5 (30-60)</td> <td>0.7-1.3 (20-40)</td> </tr> </tbody> </table> |                                   | Parenteral Ca<br>mmol (mg)/kg/day | Parenteral Ph<br>mmol (mg)/kg/day | Preterm during the first days of life | 0.8-2.0 (32-80) | 1.0-2.0 (31-62) | Growing preterm | 1.6-3.5 (100-140) | 1.6-3.5 (77-108) | Term neonate | 0.8-1.5 (30-60) | 0.7-1.3 (20-40) |
|                                       | Parenteral Ca<br>mmol (mg)/kg/day  | Parenteral Ph<br>mmol (mg)/kg/day |                                   |                                   |                                       |                 |                 |                 |                   |                  |              |                 |                 |
| Preterm during the first days of life | 0.8-2.0 (32-80)  | 1.0-2.0 (31-62)                   |                                   |                                   |                                       |                 |                 |                 |                   |                  |              |                 |                 |
| Growing preterm                       | 1.6-3.5 (100-140)  | 1.6-3.5 (77-108)                  |                                   |                                   |                                       |                 |                 |                 |                   |                  |              |                 |                 |
| Term neonate                          | 0.8-1.5 (30-60)  | 0.7-1.3 (20-40)                   |                                   |                                   |                                       |                 |                 |                 |                   |                  |              |                 |                 |

**Enteral intake:** ESPGHAN 2010 Guidelines for enteral nutrition recommend 2–3 mmol/kg/day of a highly absorbable phosphate source in a ratio with calcium (Ca:P) of 1.5–2.0.<sup>7</sup> American Academy of Pediatrics Committee on Nutrition 2013 Guidelines recommend Ca 150-200 mg/kg/day (3.8-5 mmol/kg/day) and P 75-140 mg/kg/day (2.4-4.5 mmol/kg/day) and 200-400 IU/day of vitamin D for enteral nutrition in preterm neonates.<sup>8</sup>

The exact serum phosphorus concentration at which to commence supplementation of phosphate is not known and recommendations vary from 1.3 mmol/L<sup>8</sup> to 1.8 mmol/L.<sup>9</sup>

### Metabolic bone disease

Goal: Aim for the upper end of the recommended range to prevent fractures and clinical symptoms of osteopenia: Ca and P of around 4-4.5 mmol/kg/day. Adjust the mineral intake with a goal of achieving a slight excess of urinary mineral excretion: Urinary calcium  $\geq$ 1.2mmol/L and phosphate  $\geq$ 0.4 mmol/L.<sup>14</sup>

### Step 1: Calculate the mineral intake from enteral feed:

Example: 150 ml/kg/day of mature preterm EBM contains: Ca 1 mmol/kg/day and P 0.6 mmol/kg/day.  
150 ml/kg/day preterm EBM+24kcal HMF contains: Ca 4.5 mmol/kg/day and P 2.7 mmol/kg/day.

| Preterm milk         | Ca, mmol (mg)/100 mL | P, mmol (mg)/100 mL |
|----------------------|----------------------|---------------------|
| 1 <sup>st</sup> week | 0.7 (26)             | 0.4 (11)            |
| 2 <sup>nd</sup> week | 0.6 (25)             | 0.5 (15)            |
| Week 3/4             | 0.6 (25)             | 0.5 (14)            |
| Week 10/12           | 0.7 (29)             | 0.4 (12)            |
| Term milk            |                      |                     |
| 1 <sup>st</sup> week | 0.7 (26)             | 0.4 (12)            |
| 2 <sup>nd</sup> week | 0.7 (28)             | 0.6 (17)            |
| Week 3/4             | 0.7 (27)             | 0.5 (16)            |
| Week 10/12           | 0.7 (26)             | 0.5 (16)            |

Elemental Ca, 1 mmol = 40 mg. Elemental Phosphorus, 1 mmol = 31 mg. Adapted from Gidrewicz and Fenton BMC Pediatrics 2014, 14:216.<sup>15</sup>

### Step 2: Calculate the gap in Ca and P intake/requirement: This will be the dose required.

**Step 3: Prescribe 50% of the required dose of Ca and P in 2-3 divided doses alternatively but not together.** (example: Ca 8 AM, 2 PM, 8 PM and P 6 AM, 12 MD, 6 PM).

### Step 4: Once 50% dose is tolerated for 1 week, increase to 100% required dose.

ORAL preparation during NICU stay: Sodium dihydrogen phosphate Phebra IV is the preferred preparation for oral administration due to its low osmolality.

ORAL preparation at discharge or stable neonates: Phosphate effervescent tablets can be used.

American Academy of Pediatrics Committee on nutrition 2013 Guidelines on management for Enterally Fed Preterm Infants With Radiologic Evidence of Rickets: 1. Maximize nutrient intake. 2. If no further increases in these can be made, add elemental calcium and phosphorus as tolerated. Usually beginning at 20 mg/kg per day of elemental calcium and 10–20 mg/kg per day elemental phosphorus and increasing, as tolerated, usually to a maximum of 70–80 mg/kg per day of elemental calcium and 40–50 mg/kg per day elemental phosphorus. May consider targeting 25-OH-D concentration of  $>$ 20 ng/mL (50 nmol/L).<sup>8</sup> However, breast milk content of phosphorus is variable and harder to estimate the intakes accurately. A more pragmatic approach suggested by our consensus group: start with P 0.5-1.0 mmol/kg/day in divided doses and increase as tolerated to a maximum of P 3 mmol/kg/day.

### Efficacy and safety

An ideal oral form of phosphate for use in preterm infants does not exist. Administering the intravenous preparations orally can be considered, because they are lower in osmolality than are commercially available phosphorus-containing liquids. For example, potassium dihydrogen phosphate provides 31 mg

|                        |   |
|------------------------|---|
|                        | <p>of elemental phosphorus per millimole. A dose of 10 to 20 mg/kg per day of elemental phosphorus is reasonable and will likely resolve hypophosphataemia in most preterm infants.<sup>8</sup></p> <p>Oral phosphorus and feeds</p> <p>It is recommended to separate oral doses from calcium and antacids containing agents such as aluminium hydroxide, calcium or magnesium salts, as these may reduce the bioavailability of phosphate. Oral phosphate preparation has high osmolality and administration with feeds may have theoretical benefit of reducing the osmolality (consensus opinion).</p>   |
| <b>Practice points</b> |   |
| <b>References</b>      | <ol style="list-style-type: none"> <li>1. Tsang R, Uauy R, Koletzko B, Zlotkin SH. Calcium, magnesium, phosphate and vitamin D. In Nutrition of the preterm infant. Scientific basis and practical guidelines 2005:p 265.</li> <li>2. Dissaneewate S, Vachvanichsanong P. Severe hyperphosphatemia in a newborn with renal insufficiency because of an erroneous medical prescription. Journal of Renal Nutrition. 2009 Nov 30;19(6):500-2.</li> <li>3. Schanler RJ, Abrams SA, Garza C. Mineral balance studies in very low birth weight infants fed human milk. J Pediatr. 1988;113 (1 pt 2):230–238.</li> <li>4. AAP. Pediatric nutrition handbook. 6th ed. Kleinman RE, editor: AAP eBooks; 2009.</li> <li>5. Koletzko B, Goulet O, Hunt J, Krohn K, Shamir R. 1. Guidelines on Paediatric Parenteral Nutrition of the European Society of Paediatric Gastroenterology, Hepatology and Nutrition (ESPGHAN) and the European Society for Clinical Nutrition and Metabolism (ESPEN), Supported by the European Society of Paediatric Research (ESPR). J Pediatr Gastroenterol Nutr. 2005;41 Suppl 2:S1-87.</li> <li>6. Bolisetty S, Osborn D, Sinn J, Lui K. Australasian Neonatal Parenteral Nutrition Consensus Group. Standardised neonatal parenteral nutrition formulations-an Australasian group consensus 2012. BMC Pediatr. 2014;14:48.</li> <li>7. Agostoni C, Buonocore G, Carnielli VP, et al; ESPGHAN Committee on Nutrition. Enteral nutrient supply for preterm infants: commentary from the European Society of Paediatric Gastroenterology, Hepatology and Nutrition Committee on Nutrition. J Pediatr Gastroenterol Nutr. 2010;50(1):85–91.</li> <li>8. Abrams SA, and the Committee on nutrition. Calcium and Vitamin D Requirements of Enterally Fed Preterm Infants. Pediatrics 2013;131:e1676–e1683.</li> <li>9. Tinnion RJ, Embleton ND. How to use... alkaline phosphatase in neonatology. Arch Dis Child Educ Pract 2012;97:157–63.</li> <li>10. Bozzetti V, Tagliabue P. Metabolic Bone Disease in preterm newborn: an update on nutritional issues. Italian J Ped 2009;35:20. Doi:10.1186/1824-7288-35-20.</li> <li>11. MIMS Product Info. Accessed on 11 April 2018.</li> <li>12. Mihatsch W, et al., ESPGHAN/ESPEN/ESPR/CSPEN guidelines on pediatric parenteral nutrition: Calcium, phosphorus and magnesium, Clinical Nutrition (2018), <a href="https://doi.org/10.1016/j.clnu.2018.06.950">https://doi.org/10.1016/j.clnu.2018.06.950</a>.</li> <li>13. Schanler RJ, Atkinson SA. Human milk. In Nutrition of the preterm infant. Scientific basis and practical guidelines. Second edition 2005. Eds Tsang R, Uauy R, Koletzko B, Zlotkin SH.:p 336.</li> <li>14. Osborn DA. Metabolic bone disease. <a href="https://www.slhd.nsw.gov.au/rpa/neonatal%5Ccontent/pdf/guidelines/metabolicBD.pdf">https://www.slhd.nsw.gov.au/rpa/neonatal%5Ccontent/pdf/guidelines/metabolicBD.pdf</a></li> <li>15. Gidrewicz DA, Fenton TR. A systematic review and meta-analysis of the nutrient content of preterm and term breast milk. BMC pediatrics. 2014 Dec;14(1):216.</li> </ol> |

| VERSION/NUMBER | DATE       |
|----------------|------------|
| Original 1.0   | 15/11/2016 |
| Version 1.1    | 22/11/2018 |
| Version 2.0    | 20/05/2021 |
| Version 3.0    | 15/07/2021 |
| Current 4.0    | 21/10/2021 |
| REVIEW         | 21/10/2026 |

### Authors Contribution

|                   |                                |
|-------------------|--------------------------------|
| Original author/s | Chris Wake, Srinivas Bolisetty |
| Expert review     |                                |

|  |   |
|--|---|
| Current version author                           | Srinivas Bolisetty  |
| Evidence Review                                  | David Osborn  |
| Nursing Review                                   | Eszter Jozsa, Priya Govindaswamy  |
| Pharmacy Review                                  | Jing Xiao, Cindy Chen   |
| ANMF Group contributors                          | Nilkant Phad, Bhavesh Mehta, John Sinn, Michelle Jenkins, Thao Tran, Helen Huynh, Simarjit Kaur, Jessica Mehegan, Joanne Malloy, Mohammad Irfan Azeem |
| Final content and editing review of the original | Ian Whyte   |
| Electronic version                               | Cindy Chen, Ian Callander   |
| Facilitator                                      | Srinivas Bolisetty  |