

Indometacin (Indomethacin)

Newborn use only

2021

Alert	From April 2016, the international spelling for Indomethacin has been changed to Indometacin.			
Indication	Closure of patent ductus arteriosus (PDA) Prevention of severe intra-ventricular haemorrhage.			
Action	Prostaglandin inhibitor. Prostaglandins are important in maintaining ductal patency in utero.			
Drug type	Non-steroidal anti-inflammatory drug (NSAID).			
Trade name	Indocid PDA, Indomethacin Agila			
Presentation	1 mg powder for reconstitution.			
Dose	IV Single daily dose as follows:			
	Post-natal Age	Day 1	Day 2	Day 3
	< 48 hours	0.2 mg/kg/dose	0.1 mg/kg/dose	0.1 mg/kg/dose
	≥ 48 hours	0.2 mg/kg/dose	0.2 mg/kg/dose	0.2 mg/kg/dose
Dose adjustment	Not applicable			
Therapeutic hypothermia	Insufficient data to suggest dose adjustments.			
ECMO	Refer to contraindications section.			
Renal impairment	Insufficient data to suggest dose adjustments.			
Hepatic impairment				
Maximum dose	0.2 mg/kg			
Total cumulative dose	0.6 mg/kg			
Route	IV			
Preparation	Add 1 mL of WFI to the 1 mg powder for reconstitution. Then draw up 1 mL (1 mg) and add 9 mL WFI to make a final volume of 10 mL with a concentration of 0.1 mg/mL.			
Administration	IV: Over 20–30 minutes. Inspect visually for particulate matter and discolouration prior to administration.			
Monitoring	Monitor urine output, cardiovascular status, serum biochemistry, renal function and for signs of bleeding.			
Contraindications	Renal impairment: urine output <1 mL/kg/hour during the preceding 8 hours; serum creatinine ≥140 µmol/L; blood urea nitrogen >14 mmol/L. (17) Serious infection, active bleeding, thrombocytopenia or coagulopathy, necrotising enterocolitis (NEC) or intestinal perforation, significant renal dysfunction, ductal dependent congenital heart disease and pulmonary hypertension.			
Precautions	Indomethacin is associated with transient renal impairment. Late and prolonged treatment of the ductus arteriosus with indomethacin may increase the incidence of NEC.			
Drug interactions	Aminoglycosides: Dose may need to be modified if indomethacin affects renal function. Digoxin: Reduces indomethacin volume of distribution – increased dose may be required. Diuretics: Use of frusemide in combination with indomethacin may increase the incidence of renal impairment. Systemic corticosteroids: Intestinal perforation has been described in infants treated with early dexamethasone and indomethacin.			
Adverse reactions	Prophylactic indomethacin is associated with oliguria/anuria. Treatment of the ductus arteriosus with indomethacin and prolonged courses of indomethacin are associated with NEC. Gastrointestinal perforation and possibly bleeding. Extravasation.			
Compatibility	Fluids: Sodium chloride 0.9%, water for injection. Y site: Atropine, Cephazolin, cefotaxime, ceftazidime, clindamycin, dexamethasone, digoxin, fentanyl, fluconazole, frusemide, heparin, hydrocortisone, benzylpenicillin, potassium chloride, sodium bicarbonate.			
Incompatibility	Fluids: Glucose 7.5%, Glucose 10% Y-site: Amino acid solutions, adrenaline, amikacin, atracurium, aztreonam, buprenorphine, buprenorphine, calcium chloride, calcium gluconate, chlorpromazine, dobutamine, dopamine, erythromycin, esmolol, gentamicin, glycopyrrolate, haloperidol lactate, hydralazine, labetalol, magnesium sulfate, metaraminol,			

Indometacin (Indomethacin)

Newborn use only

2021

	midazolam, morphine sulfate, noradrenaline, ondansetron, pentamidine, pethidine, phenylephrine, promethazine, protamine, suxamethonium, tobramycin, vancomycin, vasopressin, verapamil.
Stability	Discard unused portion. Diluted solution is stable for 6 hours at room temperature.
Storage	Store unopened vials at room temperature (20–25°C)
Excipients	
Special comments	
Evidence	<p>Effectiveness: Prophylactic intravenous indomethacin in preterm infants has short-term benefits including a reduction in the incidence of symptomatic PDA, PDA surgical ligation and severe intraventricular haemorrhage (IVH). However, there is no evidence of effect on mortality or neurodevelopment (5) (LOE I GOR C). Safety: Prophylactic indomethacin is associated with oliguria but not an increased creatinine or gastrointestinal side effects.</p> <p>Indomethacin for asymptomatic patent ductus arteriosus: Treatment of an asymptomatic PDA with indomethacin reduced the incidence of symptomatic PDA, duration of supplemental oxygen, with no effect on mortality, IVH, retinopathy of prematurity, length of ventilation, or NEC. Safety: Renal and gastrointestinal toxicities and long term neurodevelopment were not reported (10) (LOE I, GOR C).</p> <p>Indomethacin versus ibuprofen for the treatment of patent ductus arteriosus in preterm or low birth weight infants: Indomethacin is as effective as ibuprofen in closing a PDA (6). Safety: Indomethacin increases the risk of NEC and transient renal insufficiency compared to ibuprofen.</p> <p>Summary recommendation: Ibuprofen is as effective as indomethacin in closing a PDA and currently appears to be the drug of choice. Ibuprofen reduces the risk of NEC and transient renal insufficiency compared to indomethacin (6) (LOE I GOR B).</p> <p>Dose: Indomethacin given in total amounts for the prolonged course (6–8 doses) of 0.6–1.6 mg/kg compared with the short course 0.3–0.6 mg/kg (2–3 doses): There was no difference in efficacy between a short or prolonged course of indomethacin (LOE 1, GOR C). Safety: A prolonged course is associated with an increased risk of NEC but a decreased incidence of renal function impairment (oliguria and increased serum creatinine) (7) (LOE I, GOR B). Pharmacokinetic studies reported substantial interpatient variability (11, 12) in clearance related to postnatal age. (2, 12) Bolus infusions of indomethacin are associated with alterations in renal, mesenteric and cerebral blood flow (13). Ductus arteriosus closure rates are related to dose and indomethacin concentrations.(11,14).</p>
Practice points	
References	<ol style="list-style-type: none"> Allegaert K. The impact of ibuprofen or indomethacin on renal drug clearance in neonates. <i>The journal of maternal-fetal & neonatal medicine</i>. 2009;22:88–91. Smyth JM, Collier PS, Darwish M, Millership JS, Halliday HL, Petersen S, McElnay JC. Intravenous indomethacin in preterm infants with symptomatic patent ductus arteriosus. A population pharmacokinetic study. <i>British journal of clinical pharmacology</i>. 2004;58:249–58. Lee BS, Byun SY, Chung ML, Chang JY, Kim HY, Kim EA, Kim KS, Pi SY. Effect of furosemide on ductal closure and renal function in indomethacin-treated preterm infants during the early neonatal period. <i>Neonatology</i>. 2010;98:191–9. Brion LP, Campbell DE. Furosemide for symptomatic patent ductus arteriosus in indomethacin-treated infants. <i>The Cochrane database of systematic reviews</i>. 2001:CD001148. Fowlie PW, Davis PG, McGuire W. Prophylactic intravenous indomethacin for preventing mortality and morbidity in preterm infants. <i>The Cochrane database of systematic reviews</i>. 2010:CD000174. Ohlsson A, Walia R, Shah SS. Ibuprofen for the treatment of patent ductus arteriosus in preterm or low birth weight (or both) infants. <i>The Cochrane database of systematic reviews</i>. 2015;2:CD003481. Herrera C, Holberton J, Davis P. Prolonged versus short course of indomethacin for the treatment of patent ductus arteriosus in preterm infants. <i>The Cochrane database of systematic reviews</i>. 2007:CD003480. Stark AR, Carlo WA, Tyson JE, Papile LA, Wright LL, Shankaran S, Donovan EF, Oh W, Bauer CR, Saha S, Poole WK, Stoll BJ, National Institute of Child H, Human Development Neonatal Research N. Adverse effects of early dexamethasone in extremely-low-birth-weight infants. <i>National Institute of Child Health</i>

Indometacin (Indomethacin)

Newborn use only

2021

and Human Development Neonatal Research Network. The New England journal of medicine. 2001;344:95–101.

9. Walker SE, Gray S, Schmidt B. Stability of reconstituted indomethacin sodium trihydrate in original vials and polypropylene syringes. American journal of health-system pharmacy : AJHP : official journal of the American Society of Health-System Pharmacists. 1998; 55:154–8.

10. Cooke L, Steer P, Woodgate P. Indomethacin for asymptomatic patent ductus arteriosus in preterm infants. The Cochrane database of systematic reviews. 2003:CD003745.

11. Brash AR, Hickey DE, Graham TP, Stahlman MT, Oates JA, Cotton RB. Pharmacokinetics of indomethacin in the neonate. Relation of plasma indomethacin levels to response of the ductus arteriosus. The New England journal of medicine. 1981; 305:67–72.

12. Yaffe SJ, Friedman WF, Rogers D, Lang P, Ragni M, Sacchar C. The disposition of indomethacin in preterm babies. The Journal of pediatrics. 1980;97:1001–6.

13. Gork AS, Ehrenkranz RA, Bracken MB. Continuous infusion versus intermittent bolus doses of indomethacin for patent ductus arteriosus closure in symptomatic preterm infants. The Cochrane database of systematic reviews. 2008:CD006071.

14. Shaffer CL, Gal P, Ransom JL, Carlos RQ, Smith MS, Davey AM, Dimaguila MA, Brown YL, Schall SA. Effect of age and birth weight on indomethacin pharmacodynamics in neonates treated for patent ductus arteriosus. Critical care medicine. 2002;30:343–8.

15. Indomethacin. In: IV index. Trissel's 2 clinical pharmaceuticals database (parenteral compatibility). Greenwood Village, Colorado: Truven Health Analytics. Accessed 9/9/15.

16. Australian Injectable Drugs Handbook, 6th Edition, Society of Hospital Pharmacists of Australia 2014.

17. Van Overmeire B, Smets K, et al. A comparison of ibuprofen and indomethacin for closure of patent ductus arteriosus. N Engl J Med. 2000 Sep 7; 343(10):674-81.

VERSION/NUMBER	DATE
Original	29/09/2015
Version 1.2	25/12/2016
Version 1.3	20/07/2020
Version 2.0	18/02/2021
Current 3.0	23/09/2021
REVIEW	23/09/2026

Authors Contribution

Original author/s	David Osborn, Srinivas Bolisetty
Evidence Review	David Osborn, Tim Schindler
Expert review	
Nursing Review	Eszter Jozsa, Jan Polverino
Pharmacy Review	Mariella De Rosa, Ushma Trivedi
ANMF Group contributors	Himanshu Popat, Nilkant Phad, Michelle Jenkins, Thao Tran, Carmen Burman, Cindy Chen, Wendy Huynh, John Sinn, Renae Gengaroli, Simarjit Kaur
Final editing and review of the original	Ian Whyte
Electronic version	Cindy Chen, Ian Callander
Facilitator	Srinivas Bolisetty